Developing a Laboratory Test to Diagnose and Subtype Autism using Metabonomics

	1					
In	sti		141	\sim	n	
	э.	ıı	JLI	u		_

University of South Australia (SA) and Flinders University (SA)

Tags:

Laboratory Test, Diagnose Subtype, Metabonomics

Download summary:

External links:

Citation:

Wang LV, Angley MT, Sorich MJ, Young RL, McKinnon RA & Gerber JP, (2010) Is there a role for routinely screening children with autism spectrum disorder for creatine deficiency syndrome? Autism Research. Vol.3 pp. 268–272, 2010 DOI: 10.1002/aur.145.

Yap IKS, Angley M, Veselkov K, Holmes E, Lindon JC, Nicholson JK. (2010) Urinary metabolic phenotyping differentiates children with autism, from their unaffected siblings and age-matched controls. Journal of Proteome Research Vol. 9 pp. 2996–3004 DOI: 10.1021/pr901188e

Wang LV, Angley MT, Gerber JP, Young RL, Abarno DV, McKinnon RA & Sorich MJ, (2009) Is Urinary Indolyl-3-Acryloylglycine (IAG) a Biomarker for Autism with Gastrointestinal Symptoms? Biomarkers. Vol: 14, pp 596-603 DOI 10.1080/13547500903183962

Wang LV, Christophersen CT, Sorich MJ, Gerber, CP, Angley MT, Conlon MA, (2011) Low Relative Abundances of the Mucolytic Bacterium Akkermansia muciniphila and Bifidobacterium spp. in Feces of Children with Autism. Applied and Environmental Microbiology Vol. 77 (18) pp. p. 6718–6721, doi:10.1128/AEM.05212-11v [IF: 3.6].

Project ID:

MA200502